



Complement MASP-1 enhances adhesion between endothelial cells and neutrophils by up-regulating E-selectin expression



Péter K. Jani^{a,1}, Endre Schwaner^{a,1}, Erika Kajdáczi^a, Márta L. Debreczeni^a, Rita Ungai-Salánki^{b,c,d}, József Dobó^e, Zoltán Doleschall^f, János Rigó Jr.^g, Miklós Geiszt^h, Bálint Szabó^{c,d}, Péter Gál^e, László Cervenak^{a,*}

^a 3rd Department of Internal Medicine, Semmelweis University, 1125 Budapest, Hungary

^b Doctoral School of Molecular and Nanotechnologies, University of Pannonia, 8200 Veszprém, Hungary

^c Nanobiosensorics Lendület Group, Institute for Technical Physics and Materials Science, Centre for Energy Research, 1121 Budapest, Hungary

^d Department of Biological Physics, Eötvös Loránd University, 1053 Budapest, Hungary

^e Institute of Enzymology, Research Centre for Natural Sciences, Hungarian Academy of Sciences, 1117 Budapest, Hungary

^f Department of Pathogenetics, National Institute of Oncology, 1122 Budapest, Hungary

^g First Department of Obstetrics and Gynecology, Semmelweis University, 1088 Budapest, Hungary

^h Department of Physiology, Semmelweis University, 1094 Budapest, Hungary

ARTICLE INFO

Article history:

Received 16 March 2016

Received in revised form 5 May 2016

Accepted 7 May 2016

Keywords:

Adhesion
Complement
Endothelial cell
Neutrophil
Inflammation
MASP-1

ABSTRACT

The complement system and neutrophil granulocytes are indispensable in the immune response against extracellular pathogens such as bacteria and fungi. Endothelial cells also participate in antimicrobial immunity largely by regulating the homing of leukocytes through their cytokine production and their pattern of cell surface adhesion molecules. We have previously shown that mannan-binding lectin-associated serine protease-1 (MASP-1), a complement lectin pathway enzyme, is able to activate endothelial cells by cleaving protease activated receptors, which leads to cytokine production and enables neutrophil chemotaxis. Therefore, we aimed to investigate how recombinant MASP-1 (rMASP-1) can modify the pattern of P-selectin, E-selectin, ICAM-1, ICAM-2, and VCAM-1 adhesion molecules in human umbilical vein endothelial cells (HUVEC), and whether these changes can enhance the adherence between endothelial cells and neutrophil granulocyte model cells (differentiated PLB-985). We found that HUVECs activated by rMASP-1 decreased the expression of ICAM-2 and increased that of E-selectin, whereas ICAM-1, VCAM-1 and P-selectin expression remained unchanged. Furthermore, these changes resulted in increased adherence between differentiated PLB-985 cells and endothelial cells. Our finding suggests that complement MASP-1 can increase adhesion between neutrophils and endothelial cells in a direct fashion. This is in agreement with our previous finding that MASP-1 increases the production of pro-inflammatory cytokines (such as IL-6 and IL-8) and chemotaxis, and may thereby boost neutrophil functions. This newly described cooperation between complement lectin pathway and neutrophils via endothelial cells may be an effective tool to enhance the antimicrobial immune response.

© 2016 Elsevier Ltd. All rights reserved.

1. Introduction

The effectiveness of the immune system depends on the prompt and specific response against pathogens. Therefore, it is not

Abbreviation: HUVEC, human umbilical vein endothelial cell; MBL, mannan-binding lectin; MASP, mannan-binding lectin-associated serine protease; PAR, protease-activated receptor.

* Corresponding author at: 3rd Department of Internal Medicine, Semmelweis University, Kútvolgyi street 4, 1125 Budapest, Hungary.

E-mail addresses: cervenak.laszlo@med.semmelweis-univ.hu, cervenak.laszlo@gmail.com (L. Cervenak).

¹ These authors contributed equally to this work.

<http://dx.doi.org/10.1016/j.molimm.2016.05.007>

0161-5890/© 2016 Elsevier Ltd. All rights reserved.

surprising that immunological subsystems cooperate with one another for the quick recognition and elimination of invaders or altered host cells. A good example of this cooperation is the teamwork of the complement system and neutrophil granulocytes against bacteria and fungi.

The complement system can be activated through different routes: the classical, the alternative, and the lectin pathways. After recognition of the target structure by any of the three pathways, a cascade of serine proteases is activated, and this leads to the cleavage of C3, the central molecule of the complement system. C3b, C4b and their degradation fragments act as opsonins enhancing the phagocytic activity of neutrophils (and other cell types).

C3a and C5a are amongst the most potent chemotactic factors for neutrophils attracting them to the site of the infection.

Mannan-binding lectin (MBL)-associated serine protease-1 (MASP-1) is the primary enzyme of the lectin pathway (Heja et al., 2012a; Heja et al., 2012b; Megyeri et al., 2013). MASP-1 is activated upon the recognition of special pathogen/danger-associated motifs by MBL, collectin-11 (CL-K1), or ficolins (1-, 2-, and 3-ficolin) (Kjaer et al., 2013; Matsushita et al., 2013). Following its autoactivation, MASP-1 cleaves MASP-2 and C2, thereby ensuring further activation of the lectin pathway. However, we and other groups have described several other functions of MASP-1 beyond complement activation (Dobo et al., 2014). MASP-1 can cleave high-molecular-weight kininogen to bradykinin, a vasoactive and nociceptive peptide (Dobo et al., 2011). It can also cleave prothrombin, FXIII and TAFI, thus MASP-1 can modulate the function of the clotting and fibrinolytic systems (Jenny et al., 2015; Krarup et al., 2008; Hess et al., 2012). Moreover, active MASP-1 (both in native form complexed with MBL and as recombinant, rMASP-1) is able to cleave protease activated receptors (PAR1, 2 and 4), and this generates a pro-inflammatory signal in endothelial cells (Megyeri et al., 2009; Megyeri et al., 2014). Upon this stimulus, the endothelial cells produce IL-6 and IL-8 and thus, recruit neutrophil granulocytes (Jani et al., 2014).

Endothelial cells are vascular site specific regulators of several physiological processes (Aird, 2007a; Aird, 2007b). To interact with circulating blood cells, endothelial cells must utilize a wide range of adhesion molecules. These adhesion molecules can be classified according to their structure, subcellular localization, and function. Upon traumatic vascular injury, endothelial cells translocate their pre-formed P-selectin molecules onto the cell surface by degranulation occurring within a few minutes (Hattori et al., 1989; Rondajaj et al., 2006). P-selectin is the major lectin-type adhesion molecule of endothelial cells that interacts with platelets to ensure their adherence to the injured vessel wall. Inflammation induced by invading microorganisms or necrotic debris stimulates endothelial cells to modify the pattern of other adhesion molecules. Sialyl-Lewis X recognizing E-selectin is *de novo* synthesized by endothelial cells within hours of induction to enhance the initial adhesion and rolling of leukocytes on the endothelium (Abbassi et al., 1993; Walz et al., 1990). ICAM-1, a receptor of beta-2 integrins, is expressed on the surface of most endothelial cells (and of some leukocytes) at a low level; however, its expression is up-regulated within 12–24 h in response to pro-inflammatory stimuli (Tonnesen, 1989; Dustin et al., 1986; Poher et al., 1987). It participates in the transmigration of all types of leukocytes. VCAM-1 is another inflammatory adhesion molecule synthesized *de novo*, but only later (24 h after) an appropriate stimulus. As a partner of VLA-4, it has a major role in the homing of T-cells and monocytes, as well as subsets of neutrophils also use this adhesion molecule (Carlos and Harlan, 1994; Elices et al., 1990). ICAM-2 is an endothelial cell specific adhesion molecule. Although it is constitutively expressed on non-activated endothelial cells, its exact role is still not fully understood. ICAM-1 and ICAM-2 share their integrin ligands. However, the affinity spectrum of ICAMs towards different beta-2 integrins differ, and ICAM-2 does not induce outside-in signaling of integrin partner molecules (Shimaoka et al., 2001; Thompson et al., 2002). ICAM-2 expression decreases upon pro-inflammatory stimuli (McLaughlin et al., 1998).

To control leukocyte homing during inflammation, the production of cytokines and the expression of adhesion molecules are usually co-regulated in endothelial cells. We have previously described that rMASP-1 can induce IL-6 and IL-8 production in HUVECs, which is regulated predominantly by the p38-MAPK pathway (Jani et al., 2014). Therefore, in this study we investigated how rMASP-1 can modify the pattern of P-selectin, E-selectin, ICAM-1, ICAM-2, and VCAM-1 adhesion molecules in endothelial cells. We

also explored whether these changes can enhance the adherence between endothelial cells and neutrophil granulocytes.

2. Material and methods

2.1. Reagents

We used recombinant catalytic fragment of human MASP-1 (CCP1-CCP2-SP, hereinafter: rMASP-1). rMASP-1 was expressed in *E. coli* as described by Ambrus et al. (Ambrus et al., 2003), and prepared according to Dobó et al. (Dobo et al., 2008). Briefly, after dissolving inclusion bodies and refolding in glutathione/oxidized glutathione containing buffer for one month, rMASP-1 autoactivated during dialysis. Then sequential ion-exchange chromatographies of rMASP-1 were performed using anion-exchange (Source 30Q, GE Healthcare) and cation-exchange (SP Sepharose High Performance, GE Healthcare) columns. Purity was checked by PAGE, whereas enzymatic activity was measured by SLGR-AMC substrate cleaving assay. The rMASP-1 preparations were free of bacterial contaminations, since its NF- κ B nuclear translocating activity could not be blocked with Polymyxin B or DNase, unlike control LPS and bacterial DNA. Furthermore, it could be inhibited by C1-Inhibitor as described previously (Megyeri et al., 2009; Megyeri et al., 2014; Jani et al., 2014). The mouse anti-human P- and E-selectin, ICAM-1, ICAM-2 and VCAM-1 antibodies were purchased from Bender MedSystems (Affimetrix, Inc., San Diego). The FITC-conjugated mouse anti-human LFA-1 (CD11a/CD18), MAC-1 (CD11b/CD18), CD15 and CD49d antibodies were purchased from ImmunoTools (ImmunoTools GmbH, Germany). The Alexa Fluor[®] 568-conjugated goat anti-mouse secondary antibody was purchased from Invitrogen (Invitrogen Co., Oregon, USA) and the horseradish-peroxidase (HRP) conjugated secondary antibody was obtained from Southern Biotech (Southern Biotech, Birmingham, USA). The recombinant human E-selectin was supplied by Sino Biological (Sino Biological Inc., Beijing, P.R. China). All other reagents were purchased from Sigma-Aldrich, unless otherwise stated.

2.2. Preparation and culturing of human umbilical vein endothelial cells (HUVECs)

Cells were harvested from fresh umbilical cords obtained during normal deliveries of healthy neonates by collagenase digestion as described earlier (Jani et al., 2014; Oroszlan et al., 2006). HUVECs were kept in gelatin-precoated flasks (Corning[®] Costar[®]) in MCDB131 medium (Life Technologies) completed with 5% heat-inactivated fetal calf serum (FCS), 2 ng/mL human recombinant epidermal growth factor (R&D Systems), 1 ng/mL human recombinant basic fibroblast growth factor (Sigma), 0.3% Insulin Transferrin Selenium (Life Technologies), 1% Chemically Defined Lipid Concentrate (Life Technologies), 1% Glutamax (Life Technologies), 1% Penicillin-Streptomycin antibiotics (Sigma), 5 μ g/mL Ascorbic acid (Sigma), 250 nM Hydrocortisone (Sigma), 10 mM Hepes (Sigma), and 7.5 U/mL Heparin (this completed medium is hereinafter: Comp-MCDB). Each experiment was performed on at least three independent primary HUVEC cultures from different individuals before the 4th passage. The study was conducted in conformity with the WMA Declaration of Helsinki; its protocol was approved by the Semmelweis University Institutional Review Board (permission number: TUKEB64/2008), and all participants provided their written informed consent before inclusion.

2.3. Culturing of the PLB-985 cell line, as a model for neutrophils

PLB-985 cells were grown in RPMI-1640 medium (Life Technologies) supplemented with 10% FCS, 1% PEST, and 1% Glutamax. Then, the cells were differentiated into neutrophil-like cells for 6 days in

the presence of 1.25% dimethyl sulfoxide (DMSO) (Tucker et al., 1987) (the differentiated PLB-985 cells are hereinafter referred to as dPLB-985 cells). Differentiation was checked by detecting CD11b, CD11a, CD15, CD43 and CD49d expression, and by ascertaining the capability of the cells to generate hydrogen-peroxide.

2.4. Flow cytometry analysis

4×10^6 PLB-985 or dPLB-985 cells were washed and resuspended in HBSS. To detect CD11a, CD11b, CD15, CD43 and CD49d expression, 5 μ g of appropriate FITC conjugated antibodies, or the isotype control antibodies were added to the cell suspensions (5×10^5 cells in 50 μ l PBS) and incubated for 30 min at 4 °C. The samples were then washed in 250 μ l cold PBS containing 0.1% BSA, centrifuged and resuspended/fixed in PBS containing 2% formaldehyde for 15 min. Next, the samples were washed in 250 μ l PBS, centrifuged and resuspended in 300 μ l PBS. Cell surface expression of adhesion molecules was monitored with a FACScalibur flow cytometer (Becton Dickinson, Franklin Lakes, NJ). No gates were used except FSC threshold avoiding debris and other microparticles. The FACS figures were plotted using Flowing Software 2.5 (www.flowingsoftware.com).

2.5. Detection of hydrogen peroxide production

An Amplex Ultra Red (Life Technologies) based fluorescence method was used (Donko et al., 2010) to quantitate the production of reactive oxygen species. In brief, 2×10^4 cells were activated with 100 nM phorbol 12-myristate 13-acetate (PMA) or left untreated as controls. Then, fluorescence was recorded continuously for 20 min, and expressed as arbitrary units.

2.6. mRNA analysis

Following pre-treatment by various methods, HUVECs were lysed and stored in TRI[®] reagent. Total RNA purification, reverse transcription and LightCycler[®] analysis were performed as described earlier (Jani et al., 2014; Oroszlan et al., 2006). The primers (Table 1) were designed with Primer3 (v. 0.4.0) designer from the NCBI database, and produced by Bio Basic Canada Inc. β -actin and GAPDH gene-specific primers were used as internal controls. The purity and the size of the PCR products were checked by melting curve analysis, agarose gel-electrophoresis, and sequencing.

2.7. Visualization of adhesion molecules by immunofluorescence microscopy

Confluent layers (10^4 cell/well) of HUVECs were cultured in 96-well plates (Corning[®] Costar[®]) for one day, then the cells were treated with 2 μ M of rMASP-1, 300 nM thrombin, 50 μ M histamine or 10 ng/mL TNF α in 100 μ l Comp-MCDB, or left untreated. After 5 min, 6 or 24 h, the cells were fixed in 1% formaldehyde for 15 min (for P-selectin), or in methanol-acetone (1:1) for 10 min (for other adhesion molecules). Then the cells were stained with primary anti-human antibodies as indicated (all antibodies were diluted 1:500), followed by Alexa Fluor[®]568-conjugated goat anti-mouse IgG (1:500), and Hoechst 33342 (1:50000, Molecular Probes/Invitrogen). The preparations were read using an Olympus IX-81 fluorescence inverted microscope as described previously (Jani et al., 2014; Mako et al., 2010).

2.8. Measurement of adhesion molecule expression by cell-based ELISA

Confluent layers of HUVECs were cultured in 96-well plates for 24 h. Then, HUVECs were treated with 0.2, 0.6, or 2 μ M

Table 1

Primers for the analysis of mRNAs of the adhesion molecules.

E-selectin	forward:	5'-tcaagtgtgagcaattgtgaac-3'
	reverse:	5'-attctccagaggacatactgc-3'
ICAM-1	forward:	5'-acagtcacatggcaacgac-3'
	reverse:	5'-gtcactgtctgcagtgtctct-3'
ICAM-2	forward:	5'-acagccacattcaacagcac-3'
	reverse:	5'-agatgtcacgaacaggagac-3'
VCAM-1	forward:	5'-tgaccttcacccaccattga-3'
	reverse:	5'-gcattgtcatattcacagaactgc-3'

rMASP-1, 300 nM thrombin, or 10 ng/mL TNF α for 1, 3, 6, 10 or 24 h. The cells were fixed and stained with P-selectin, E-selectin, ICAM-1, ICAM-2, or VCAM-1 antibodies for 60 min at room temperature. Then, HRP-conjugated goat anti-mouse antibody and 3,3',5,5'-Tetra Methyl Benzidine (TMB) (Life Technologies) were used to measure the expression of adhesion molecules.

2.9. Measurement of PLB-985 and dPLB-985 cell adhesion to E-selectin coated plates

96-well plates were coated with 0.2, 1, or 5 μ g/mL recombinant human E-selectin (Sino Biological Inc.), or left uncoated for 1 h at 37 °C, and then blocked with PBS containing 2% BSA for 1 h at 37 °C, in 5% CO₂ atmosphere. PLB-985 or dPLB-985 cells were incubated in HBSS containing 2 μ g/mL Oregon Green[®] 488 carboxylic acid diacetate succinimidyl ester (abbreviated as OG-488, Invitrogen) for 30 min in the dark, at 37 °C, in 5% CO₂ atmosphere. Then, the cells were centrifuged, and resuspended in HBSS for 30 min. OG-488 labeled PLB-985 or dPLB-985 cell suspension was added to the E-selectin precoated wells (10000 cells/well), and incubated for 1 h, at 37 °C, in 5% CO₂ atmosphere. Then, total number of dPLB-985 cells was determined with a fluorescence plate reader (Tecan Infinite[®] M1000 Pro), at 485 nm excitation and 525 nm emission wavelengths. Thereafter, the plate was washed twice vigorously with HBSS, and the fluorescence of bound cells was monitored again. The number of bound cells normalized with the initial number of cells was calculated after three independent measurements.

2.10. Assessing dPLB-985 cell adhesion to HUVECs

Confluent layers of HUVECs were cultured in 96-well plates and treated for 6 or 24 h with 0.2, 0.6, or 2 μ M rMASP-1, 300 nM thrombin, or 10 ng/mL TNF α . Then, 2×10^4 OG-488 labeled dPLB-985 cells/well were applied to the HUVECs, and incubated for 1 h at 37 °C, in 5% CO₂ atmosphere. Total dPLB-985 cell number was monitored by fluorescence as described before. Thereafter, the plate was washed twice vigorously with HBSS using a multichannel pipettor, and the fluorescence of bound cells was monitored again. The ratio of bound to total cell numbers was calculated after three independent measurements. In some experiments, HUVECs were pre-incubated for 30 min with the following pathway inhibitors: 25 μ M JNK inhibitor (SP600125), 2 μ M p38-MAPK inhibitor (SB203580), 1 μ M ERK 1/2 inhibitor (U0126) or, 5 μ M NF κ B inhibitor (Bay-11 7082). Furthermore in some cases OG-488-labeled dPLB-985 cells were pre-treated with 1 μ g/mL recombinant E-selectin for 30 min, at 37 °C, in 5% CO₂ atmosphere, before adding to the HUVECs, in order to assess the inhibition of adhesion.

2.11. Measurement of the adhesion force between dPLB-985 cells and endothelial cells with a computer controlled micropipette method

A computer controlled micropipette (CellSorter), mounted onto an inverted fluorescence microscope (Zeiss Axio Observer A1)

(Koranyi et al., 2013; Salanki et al., 2014), was used to quantify the adhesion force between the variously pre-treated HUVECs and dPLB-985 cells. In brief, a confluent layer of HUVECs was cultured in a 35-mm tissue culture plastic Petri dish (Greiner) for one day. Then, they were treated/not treated with 2 μ M of rMASP-1, 300 nM thrombin for 6 h, or with 10 ng/mL TNF α for 24 h. 10^5 OG-488 labeled dPLB-985 cells were placed into the Petri dish containing the endothelial cells, and incubated for 60 min at room temperature. Attached dPLB-985 cells were scanned and recognized using the CellSorter software. The adhesion force was measured with a glass micropipette with an inner diameter of 70 μ m. The micropipette was automatically positioned to 100 selected cells (diameter \sim 13 μ m) one by one, in order to try to pick them up (Fig. 5A). Repeating this cycle (i.e. the positioning and pick-up process) each time on the same path with an increased vacuum (expressed in kPa), the adhesion force of cells could be precisely measured. The number of the cells was counted before and after each cycle, and the ratio of adherent to originally selected cells was calculated.

2.12. Statistical analysis

Statistical analyses were performed using Student's *t*-test, one-way ANOVA with Tukey's post-test or a post-test for linear trends, or two-way ANOVA with Bonferroni post-test (GraphPad Prism 5.01 software, GraphPad, <http://www.graphpad.com/>). A *p*-value less than 0.05 was considered statistically significant. The data are presented as means \pm SEM unless otherwise stated.

3. Results

3.1. Screening the mRNA expression of adhesion molecules

Since pro-inflammatory adhesion molecules are usually synthesized *de novo* upon stimuli, we measured the mRNA levels of adhesion molecules by qPCR—except in the case of P-selectin, which is stored as a preformed protein in Weibel-Palade bodies.

Confluent layers of HUVECs were treated with 2 μ M rMASP-1 for 1, 2, 6 or 10 h, to assess mRNA expression. rMASP-1 greatly up-regulated the mRNA expression of E-selectin and VCAM-1 in endothelial cells, although the increment was somewhat less in the latter. Maximum effect was observed between 2 and 6 h and then, the expression of both adhesion molecules declined. The level of ICAM-2 mRNA decreased until the end of the measurement, whereas the expression of ICAM-1 mRNA did not change upon rMASP-1 stimulation (Fig. 1A).

3.2. The effect of rMASP-1 on the expression of cell adhesion molecules

The altered expression of these adhesion molecules indicates the pro-inflammatory changes occurring in endothelial cells. Moreover, the effects of known endothelial cell activators (such as thrombin, TNF α or histamine) are well characterized with regard to adhesion molecules.

Thrombin and histamine induced prompt P-selectin expression at 5 min, and reuptake by endothelial cells within 60 min; however, rMASP-1 failed to increase P-selectin expression (Fig. 1B and C).

Fluorescence microscopy showed that rMASP-1, thrombin, and TNF α induced the expression of E-selectin at 6 h (Fig. 1B). Using cell-based ELISA, we demonstrated that the effect of rMASP-1 on E-selectin expression declined by 24 h, similar to that of thrombin, but unlike to that of TNF α (Fig. 1C). rMASP-1 did not modify the expression of ICAM-1 at any time (Fig. 1B and C). By contrast, both thrombin and TNF α elevated the level of ICAM-1, although with different kinetics (Fig. 1C). rMASP-1, thrombin, and TNF α

all reduced ICAM-2 expression at 6 h, and even further at 24 h (Fig. 1, B and C). Interestingly, although rMASP-1 significantly induced the expression of VCAM-1 mRNA, it had no effect on VCAM-1 protein level (Fig. 1B and C). Thrombin behaved similar to rMASP-1 in respect of VCAM-1 protein expression; however, TNF α induced VCAM-1 at 6 h, and this effect was even more pronounced at 24 h (Fig. 1B and C).

3.3. rMASP-1 dose dependence and the kinetics of E-selectin and ICAM-2 expression

It is known that the expression kinetics of various adhesion molecules depends on the inherent properties of the adhesion molecule itself, and on those of the activator. The difference in expression kinetics can regulate selective cell adhesion. To assess the influence of rMASP-1 kinetics on the expression of adhesion molecules at protein level, HUVECs were treated with 2 μ M rMASP-1 for 1, 3, 6, 10 and 24 h and then, cell-based ELISA was performed. Similar to the kinetics seen at mRNA level (Fig. 1A), the protein expression of E-selectin was increased as early as at 3 h, but the maximum effect was observed at 6 h. E-selectin expression declined until 24 h post treatment (Fig. 2A). In contrast to E-selectin, ICAM-2 level was lower at 3 h than that of the untreated control, and decreased continuously until 24 h (Fig. 2B). Then, at the times of maximum rMASP-1 effect, we assessed dose dependence. We found that treatment with rMASP-1 induced E-selectin and reduced ICAM-2 expression in a dose-dependent manner (Fig. 2C and D).

3.4. Measurement of adhesion between dPLB-985 cells and endothelial cells

To clarify the biological importance of the previously described alterations of endothelial adhesion molecules, we assessed whether these molecules can bind to neutrophil granulocytes. In these adhesion measurements, we used the acute myeloid leukemia cell line PLB-985, differentiated with DMSO toward neutrophil-like cells (dPLB-985). To verify the effectiveness of the differentiation of dPLB-985 cells, we tested the cell surface expression of adhesion molecules, and the generation of reactive oxygen species. dPLB-985 cells expressed CD11a and CD11b in increased quantities, and reduced amount of CD49d compared to non-differentiated PLB-985 cells. The expression of CD15 and CD43 did not change significantly (Fig. 3A). Moreover, dPLB-985 cells, unlike their ancestors, generated reactive oxygen species in response to 100 nM PMA, as measured by H₂O₂ production (Fig. 3B).

To test the adhesive capacity of dPLB-985 cells to E-selectin, microwell plates were coated with 0.2, 1, or 5 μ g/mL recombinant E-selectin, or left untreated. Then, OG-488-labeled PLB-985 or dPLB-985 cells were added, incubated for 1 h, washed, and the number of adherent cells was calculated. The adhesion of dPLB-985 cells to the coated plate was dose-dependent, and treatment with E-selectin in a concentration as low as 5 μ g/mL resulted in significant adhesion compared to the uncoated controls. We did not observe similar dose-dependent adhesion of non-differentiated PLB cells to E-selectin coated plates (Fig. 4A).

Since we demonstrated that rMASP-1 treatment induces E-selectin expression in HUVECs, and dPLB-985 cells can adhere to recombinant E-selectin, we assessed the adherence between dPLB-985 cells and HUVECs treated with rMASP-1. To this end, HUVECs were treated with 0.2, 0.6, or 2 μ M rMASP-1, 300 nM thrombin or 10 ng/mL TNF α for 6 or 24 h, or left untreated. Then OG-488 labeled dPLB-985 cells were applied onto the HUVECs and incubated for 1 h. The number of adhered dPLB-985 cells was measured by fluorescence, and normalized with baseline total cell number. We observed that rMASP-1 treatment increased dPLB-985 cell adhesion to HUVECs in a dose dependent manner at 6 h,




Fig. 1. Adhesion molecule patterns induced by rMASP-1 on HUVECs. (A) HUVECs were treated with 2 μ M rMASP-1 for 1, 2, 6, or 10 h. Then, the cells were lysed and stored in TRI-reagent, total mRNA was purified, and qPCR performed. Values of the quantities of the adhesion molecule mRNAs in three independent experiments were normalized for that of GAPDH, used as internal control. P-selectin was not analyzed by qPCR, because of its instant secretion from pre-formed granules. (B) HUVECs were treated with 2 μ M rMASP-1, 300 nM thrombin, 50 μ M histamine, or 10 ng/mL TNF α for 5 min (P-selectin), or for 6 h (in the case of all other adhesion molecules). Then, cells were fixed and stained with mouse monoclonal anti-adhesion molecule antibodies, followed by goat-anti-mouse Alexa 568 (red) and Hoechst 33342 (blue) as nuclear staining. Each subfigure of panel B is a representative photo of three independent experiments. (C) HUVECs were treated in 96-well plates as described in panel (B), and cellular ELISA was performed. The values represent the ratio of the optical density of treated vs. non-treated wells, and the difference was analyzed by ANOVA with Tukey's post-test. Dotted lines represent the untreated control levels. The values were calculated as the mean (\pm SEM) of three independent experiments. *: $p < 0.05$, **: $p < 0.01$, ***: $p < 0.001$. (For interpretation of the references to colour in this figure legend, the reader is referred to the web version of this article.)

but not at 24 h (Fig. 4B). Both the kinetics and the volume of this effect were similar for rMASP-1 and for thrombin. In contrast, TNF α induced enhanced adhesion to HUVECs at 6 h, but adherence was even greater by 24 h (Fig. 4C). The adhesion between dPLB-985 cells and rMASP-1-induced HUVECs could be reduced to the level of untreated HUVECs by pre-incubating the dPLB-985 cells with 1 μ g/mL soluble recombinant E-selectin for 30 min (Fig. 4D).

Expression of adhesion molecules can be triggered through various signaling pathways. We reported that most of these pathways can also be activated by rMASP-1 in HUVECs (Jani et al., 2014). We used commercially available signaling pathway inhibitors to identify the most important pathways required for the E-selectin mediated adhesion between dPLB-985 cells and rMASP-1-induced HUVECs. HUVECs had been pre-incubated with the inhibitors for 30 min and then, treated with 2 μ M rMASP-1 for 6 h. Only the p38-MAPK inhibitor was able to block this adhesion, whereas JNK,

NF κ B, and ERK 1/2 pathway inhibitors had no blocking effect (Fig. 4D).

We used a computer controlled micropipette method to quantify the adhesion force between dPLB-985 cells and rMASP-1-treated HUVECs (Kornyei et al., 2013; Salanki et al., 2014). HUVECs were induced with 2 μ M of rMASP-1, 300 nM thrombin for 6 h, or 10 ng/mL TNF α for 24 h or left untreated. Then, OG-488-labeled dPLB-985 cells were added and incubated for 1 h. The attached dPLB-985 cells were probed (i.e., by trying to pick them up) with the automated micropipette. The hydrodynamic lifting force, acting on the targeted single cells, was increased in each subsequent cycle of the measurement (Fig. 5A). More dPLB-985 cells stayed attached to rMASP-1-treated endothelial cells than to non-treated controls at higher detaching forces. We observed the same effect after thrombin treatment (Fig. 5B). Induction with TNF α for 24 h, used as positive control, was the most potent in enhancing adhesion (by 4.28-fold, ± 1.63 , $p < 0.01$ at 19.2 kPa vacuum).




Fig. 2. Dose- and time- dependence of E-selectin and ICAM-2 expression induced by rMASP-1. HUVECs were treated with 2 μ M of rMASP-1 for 1, 3, 6, 10, and 24 h (A and B), or with 0, 0.2, 0.6, 2 μ M rMASP-1 for 6 h (C) and 24 h (D). Then, relative expression (compared to untreated controls of E-selectin and ICAM-2) was measured by cell-based ELISA. Dotted lines represent the untreated control levels. The values were calculated as the mean (\pm SEM) of three independent experiments. One-way ANOVA has been performed for each subfigure. In the case of kinetic measurements $p < 0.01$ for A and $p < 0.05$ for B, and of dose-dependence experiments, ANOVA ($p < 0.05$ for C and $p < 0.001$ for D) was supplemented with post-test for linear trend ($r = 0.55$, $p < 0.01$ for C and $r = -0.69$, $p < 0.001$ for D).




Fig. 3. The effects of PLB-985 cell differentiation on adhesion molecule pattern and hydrogen peroxide generation. (A) PLB-985 cells or DMSO-differentiated PLB-985 cells were suspended in HBSS. Specific mouse antibodies or isotype controls were added to the cell suspensions and incubated for 30 min, at 4 °C. After washing, samples were fixed in 2% formaldehyde for 15 min, and suspended in PBS for flow cytometry analysis. 10^4 cells were measured in each sample. RMF: Ratio of geometric mean fluorescence intensity values of labeled cells and of respective isotype controls. (B) HRP and Amplex Ultra RedTM fluorescence substrate was added to PLB-985 or dPLB-985 cells, which were then activated with 100 nM PMA or left untreated. Fluorescence was recorded every 10 s for 20 min, and expressed as arbitrary units. One representative experiment out of three is shown.




Fig. 4. The adhesion of dPLB-985 cells to E-selectin coated plates or to rMASP-1-treated endothelial cells. (A) After 96-well plates had been pre-coated with 0, 0.2, 1, or 5 $\mu\text{g/mL}$ E-selectin, OG-488-labeled PLB-985 or dPLB-985 cells were added to the wells and incubated for 1 h, at 37 °C, in 5% CO₂ atmosphere. Then, the fluorescence of bound cells was detected after washing with HBSS. The number of bound cells in three independent measurements was calculated. (B, C) HUVECs were treated with 0, 0.2, 0.6, and 2 μM rMASP-1 for 6 and 24 h (B), with 2 μM of rMASP-1, 300 nM thrombin, or 10 ng/mL TNFalpha – or left untreated – for 6, and 24 h (C), and adhesion was measured as in panel A. (D) Adhesion of OG-488-labeled dPLB-985 to HUVECs treated with 2 μM rMASP-1 for 6 h was inhibited either by pre-incubation of HUVECs with p38-MAPK, JNK, NFkB, or ERK 1/2 inhibitor for 30 min or by pre-incubation of dPLB-985 with soluble recombinant E-selectin for 30 min. Then adhesion was assessed as described above in panel A, and the inhibition is shown as the percentage of the rMASP-1 induced adhesion. The values were calculated as the mean (\pm SEM) of at least three independent experiments. Two-way ANOVA was used for panel A, whereas one-way ANOVA has been performed for panel B, C and D. Dotted lines represent the untreated control levels. *: $p < 0.05$, **: $p < 0.01$, ***: $p < 0.001$.

4. Discussion

Infection and tissue damage initiate pro-inflammatory activation of the endothelium. This modification of the endothelial cell phenotype includes altered cytokine production and adhesion molecule pattern. In the present study, we found that rMASP-1 treatment changed the adhesion molecule pattern, reduced the expression of ICAM-2, and increased that of E-selectin, whereas ICAM-1, VCAM-1 and P-selectin did not change. Furthermore, these alterations increased the adherence between neutrophil model cells (dPLB-985) and endothelial cells.

The migration of different leukocyte subsets depends, in the first place, on the pattern of chemokines and adhesion molecules produced by the endothelium. Earlier, we have demonstrated that rMASP-1 induces pro-inflammatory processes in endothelial cells. In particular, they activate Ca²⁺, NFkB, p38-MAPK, JNK, and CREB signaling pathways by cleaving PARs (Megyeri et al., 2009; Jani et al., 2014). Moreover, rMASP-1-stimulated endothelial cells produce IL-6 and IL-8, as well as recruit neutrophil granulocytes (Jani et al., 2014). Therefore, our previous and current results together

imply a novel, endothelial cell mediated pro-inflammatory role for rMASP-1.

Although MASP-1 is the most abundant protease of the lectin pathway, its isolation from human serum is a low-yield process, and the produced protein is prone to degradation during the purification. The catalytic region of MASP-1, containing the CCP1, CCP2 and SP domains, is enzymatically equivalent to the full-length molecule. Structural studies showed that only the catalytic region is accessible to the substrates present in MBL-MASP-1 complexes, since only the CCP1-CCP2-SP domains protrude from the plane of MBL (Kjaer et al., 2015). Previously, we demonstrated equivalent cellular effects of the serum purified MBL-MASP complex, and of the recombinant catalytic fragment of human MASP-1. Serum purified MASP-1 and rMASP-1 used in equimolar concentration induced the same intensity of Ca-mobilization, and both could be completely blocked with C1-inhibitor (Megyeri et al., 2014). In the current study, we observed significant effects of MASP-1 on adhesion molecules and adhesion between 0.2 and 2 μM , which is higher than the plasma concentration of MASP-1 (~143 nM). However, at the site of injury and/or infection, MASP-1 may be trapped via the




Fig. 5. Adhesion force measurement between cells using an automated micropipette. (A) Schematic representation of the hydrodynamic adhesion force measurement. OG-488 labeled dPLB-985 cells attached to the confluent layer of HUVECs were scanned and recognized by software in the microscopic images captured on a motorized inverted microscope. A glass micropipette was automatically positioned to each detected cell one by one. Cell adhesion was probed by the application of a precisely controlled fluid flow through the micropipette. (B) Adhesion force between dPLB-985 cells and rMASP-1-treated endothelial cells. HUVECs were treated in Petri dishes with 2 μ M of rMASP-1, or 300 nM thrombin for 6 h, or with 10 ng/mL TNF α for 24 h, or left untreated. Then, OG-488-labeled dPLB-985 cells were seeded onto the HUVECs (subjected to different pre-treatments) for 1 h. The adhesion force between dPLB-985 cells and HUVECs was tested by the computer controlled micropipette method. The vacuum applied (expressed in kPa) was proportional to the hydrodynamic lifting force acting on the targeted single cells. Significant deviation from the untreated values in the high-vacuum range of the plot means that the ratio of cells exhibiting firm adhesion was increased by treatment. The values were calculated as the ratio of bound cell number of treated and non treated samples at each vacuum pressure points, and shown as the mean (\pm SEM) of at least three independent experiments. One-way ANOVA has been performed for panel B. Dotted lines represent the untreated control levels. *: $p < 0.05$, **: $p < 0.01$.

pattern recognition of MBL and ficolins, thus the local concentration may be much higher than its plasma concentration.

P-selectin, one of the best-characterized components of Weibel-Palade bodies (Hattori et al., 1989), is released rapidly in response to among others thrombin and histamine by G-protein coupled PARs (Cleator et al., 2006), and by histamine H1 receptors (Kubes and Kanwar, 1994), respectively. Although it plays a role in the inflammatory response by mediating the rolling of leukocytes on the endothelium (Kubes and Kanwar, 1994), the most critical function of P-selectin is to facilitate platelet aggregation at the sites of vascular injury (Frenette et al., 1995). In contrast to P-selectin, E-selectin is an endothelial cell-specific adhesion molecule; it recognizes sugar motifs containing sialyl-Lewis X on leukocytes, and has an important role in the initial step of the endothelium/leukocyte adhesion – leukocyte rolling (Olofsson et al., 1994; Pudielko et al., 2010). E-selectin is not detectable on the surface of normally functioning endothelial cells and it is also absent from their cytoplasm; however, upon pro-inflammatory stimuli (e.g. by TNF α , LPS, IL-1 β), it is synthesized *de novo* within hours (Kansas, 1996). Our results indicate that rMASP-1 alone possibly cannot influence the endothelial cell/platelet interaction, but it may regulate the rolling of leukocytes on endothelial cells via the upregulation of E-selectin.

ICAM-1 and ICAM-2 are members of the Ig superfamily; they bind β_2 -integrins (CD11a/CD18, CD11b/CD18 and CD11c/CD18) with different affinity (Shimaoka et al., 2001). ICAM-1 is expressed at a low level on endothelial cells with a non-inflammatory phenotype, as well as on different leukocyte subsets, and their expression is induced – similar to E-selectin – by pro-inflammatory factors (Mako et al., 2010). Unlike ICAM-1, ICAM-2 is specific to endothelial cells, as well as it is constitutively and strongly expressed on normally functioning cells. Upon pro-inflammatory stimulation, the cells reduce its expression in parallel with the elevation of ICAM-1 levels. Further, their outside-in signaling is different, as the crosslinking of ICAM-1 induces actin reorganization via RhoA. Thus, it participates in the cellular movements of endothelial cells during leukocyte transmigration, whereas ICAM-2 lacks this function (Thompson et al., 2002). VCAM-1, another endothelial

cell-specific adhesion molecule expressed upon pro-inflammatory stimuli, reaches its maximum 24 h after activation; it is a ligand for VLA-4 and other β_1 -integrins (Carlos and Harlan, 1994). VCAM-1 is required for the transmigration of T cells and monocytes; however, neutrophils may also use the VLA-4/VCAM-1 adhesion for homing (Lomakina and Waugh, 2009). rMASP-1 did not induce the expression of ICAM-1 nor VCAM-1 and thus, it may not have a direct influence on the firm contact and on the transmigration of the leukocytes through the endothelium. However, by decreasing ICAM-2, it biases the ICAM-1 to ICAM-2 ratio, and this may potentiate transmigration upon other synergistic pro-inflammatory stimuli (such as by LPS, TNF α or C5a). Moreover, the elevation of VCAM-1 expression only at mRNA level also suggests a potential for synergism. The discrepancy between mRNA and protein levels of VCAM-1 is not unknown (Pietersma et al., 1997; Volanti et al., 2004). MicroRNAs may block translation of VCAM-1 without degradation of mRNA or post-translational degradation can decrease the protein level, however, the exploration of the molecular mechanism is beyond the scope of this study.

rMASP-1 induced a unique pattern and kinetics of adhesion molecules that differs from those induced by TNF α and thrombin. The dissimilarity between rMASP-1 and TNF α is not surprising, because TNFR strongly activates NF κ B, p38-MAPK and JNK signaling pathways (Mako et al., 2010). Further, it does not trigger Ca-mobilization, whereas PARs cleaved by rMASP-1 induce Ca-mobilization, JNK and p38-MAPK phosphorylation, but only weak NF κ B nuclear translocation (Megyeri et al., 2009; Jani et al., 2014). More interestingly, the rMASP-1- and thrombin-induced patterns of adhesion molecules were also different. The expression of P-selectin and of ICAM-1 was enhanced by thrombin, but not by rMASP-1 – this can be explained by the different utilization of PARs. While thrombin cleaves PAR-1 and – to a lesser extent – PAR-4, rMASP-1 is most effective in cleaving PAR-4, followed by PAR-2 and PAR-1 (Megyeri et al., 2009). If we consider the expression profile of different PARs on endothelial cells (PAR-2 expression is the highest, followed by PAR-1, and – to a much less extent – by PAR-4) (Megyeri et al., 2009), only partial overlap can be supposed between

thrombin and rMASP-1 signaling. Differential signal transduction by the various PARs is well known and in support of our findings (Soh et al., 2010). Since we have previously found that in the case of rMASP-1, the cytokine production was regulated predominantly by p38-MAPK pathway (Jani et al., 2014), it is not unexpected that only the p38-MAPK inhibitor was able to block adhesion between dPLB-985 cells and rMASP-1-induced HUVECs.

Since rMASP-1-treated HUVECs produce IL-8 but not MCP-1 (Jani et al., 2014), and as we have found an elevated level of E-selectin, but not of ICAM-1 or of VCAM-1, we propose that rMASP-1 induced an endothelial cell phenotype most suitable for the adhesion of neutrophils. Nevertheless, we could not exclude the possibility that other adhesion molecules than E-selectin, which were not measured in this study, are also involved in the increased adhesion. HUVECs from different donors exhibit significant heterogeneity (Kiszel et al., 2007). To reduce further variances in our experiments, we used a DMSO-differentiated PLB-985 cell line—devoid of the heterogeneity of primary cells, which is a well-known and suitable cellular model to study the functional responses of neutrophils (Tucker et al., 1987; Petheo et al., 2010). Furthermore, dPLB-985 could bind to E-selectin, whereas non-differentiated PLB-985 could not, and this supports that it could be a suitable model for the assessment of adhesion. The pattern of dPLB-985 binding to HUVECs subjected to different pre-treatments was rather similar to that of the expression of E-selectin. rMASP-1 and thrombin enhanced adhesion, and elevated the E-selectin level only 6 h after treatment—TNF alpha was superior in regard of both parameters, and at any times after induction. The possible role of E-selectin in this static model of dPLB-985 binding to HUVECs is further supported by the inhibition of adhesion by soluble E-selectin. To assess the adhesion force between rMASP-1-treated HUVECs and dPLB-985 cells, we utilized the recently introduced computer controlled micropipette method (Kornyey et al., 2013; Salanki et al., 2014). Previously, it was applied to measure the adherence of macrophages and dendritic cells to fibrinogen, and to poly-L-Lys-graft-PEG. The adhesion force was in the same order of magnitude (Salanki et al., 2014) as we found in the case of HUVECs/dPLB-985 cells. Our present data suggest that rMASP-1 induced high adhesion forces between dPLB-985 cells and HUVECs. The greatest difference – compared to untreated HUVECs – was measured at the highest detaching forces. This means that the number of firmly adherent cells increased significantly.

The complement system is a powerful defense mechanism for eliminating pathogens and necrotic debris. However, one of its main functions, the opsonization of unwanted particles, requires the presence of phagocytic cells. We found that complement MASP-1 can enhance the adhesion between neutrophils and endothelial cells in a direct fashion, and may thereby accelerate the recruitment of neutrophils at the scene of infection or tissue damage. This is in agreement also with our previous finding that MASP-1 increases the production of pro-inflammatory cytokines, such as of IL-6 and IL-8. Enhanced chemotaxis and adhesion may boost neutrophil functions; however, this hypothesis should be tested by further research.

Conflict of interest

BS is a founder of CellSorter, the startup company that developed the computer controlled micropipette. The other authors declare that they have no conflicts of interest with the contents of this article.

Author contributions

PKJ and ES contributed equally to this work. PKJ, ES, EK, MLD and LC harvested and cultured HUVECs and conducted most of

the experiments. PKJ, ES and LC analyzed the results, and wrote the paper. JR organized and prepared the fresh umbilical cords obtained during normal delivery of healthy neonates for harvesting HUVECs. RUS and BS conducted and analyzed the experiments with the computer controlled micropipette. JD and PG prepared rMASP-1 and revised the manuscript. ZD conducted qPCR experiments. MG supervised Amplex Ultra Red based fluorescence method. All authors reviewed the results and approved the final version of the manuscript.

Acknowledgments

The authors thank Miklós Sándor Zoltán Kellermayer for revising the adhesion measurements at Department of Biophysics and Radiation Biology, Semmelweis University, Budapest, Hungary. We would also like to thank Róbert Horváth from the Department of Photonics, Institute of Technical Physics and Materials Science, Budapest, Hungary for facilitating the collaboration with the CellSorter startup company.

This work was supported by Grants NK100834, NK100769, NK108642, K100684 and K115623 (OTKA, Hungarian Scientific Research Fund). PG, BS and LC were supported by MedinProt Synergy project. BS was supported by the Bolyai Scholarship of the Hungarian Academy of Sciences. RUS was supported by the “Lendület” Program of the Hungarian Academy of Sciences.

References

- Abbassi, O., Kishimoto, T.K., McIntire, L.V., Anderson, D.C., Smith, C.W., 1993. E-selectin supports neutrophil rolling in vitro under conditions of flow. *J. Clin. Invest.* 92, 2719–2730.
- Aird, W.C., 2007a. Phenotypic heterogeneity of the endothelium: I. Structure, function, and mechanisms. *Circ. Res.* 100, 158–173.
- Aird, W.C., 2007b. Phenotypic heterogeneity of the endothelium: II. Representative vascular beds. *Circ. Res.* 100, 174–190.
- Ambrus, G., Gal, P., Kojima, M., Szilagyi, K., Balczar, J., Antal, J., Graf, L., Laich, A., Moffatt, B.E., Schwaeble, W., Sim, R.B., Zavodszky, P., 2003. Natural substrates and inhibitors of mannan-binding lectin-associated serine protease-1 and -2: a study on recombinant catalytic fragments. *J. Immunol.* 170, 1374–1382.
- Carlos, T.M., Harlan, J.M., 1994. Leukocyte-endothelial adhesion molecules. *Blood* 84, 2068–2101.
- Cleator, J.H., Zhu, W.Q., Vaughan, D.E., Hamm, H.E., 2006. Differential regulation of endothelial exocytosis of P-selectin and von Willebrand factor by protease-activated receptors and cAMP. *Blood* 107, 2736–2744.
- Dobo, J., Harmat, V., Sebestyen, E., Beinrohr, L., Zavodszky, P., Gal, P., 2008. Purification, crystallization and preliminary X-ray analysis of human mannan-binding lectin-associated serine protease-1 (MASP-1) catalytic region. *Acta Crystallographica. Section F. Struct. Biol. Cryst. Commun.* 64, 781–784.
- Dobo, J., Major, B., Kekesi, K.A., Szabo, I., Megyeri, M., Hajela, K., Juhasz, G., Zavodszky, P., Gal, P., 2011. Cleavage of kininogen and subsequent bradykinin release by the complement component: mannan-binding lectin-associated serine protease (MASP)-1. *PLoS One* 6, e20036.
- Dobo, J., Schroeder, V., Jenny, L., Cervenak, L., Zavodszky, P., Gal, P., 2014. Multiple roles of complement MASP-1 at the interface of innate immune response and coagulation. *Mol. Immunol.* 61, 69–78.
- Donko, A., Ruisanchez, E., Orient, A., Enyedi, B., Kapui, R., Peterfi, Z., de Deken, X., Benyo, Z., Geiszt, M., 2010. Urothelial cells produce hydrogen peroxide through the activation of Duox1. *Free Radic. Biol. Med.* 49, 2040–2048.
- Dustin, M.L., Rothlein, R., Bhan, A.K., Dinarello, C.A., Springer, T.A., 1986. Induction by IL 1 and interferon-gamma: tissue distribution, biochemistry, and function of a natural adherence molecule (ICAM-1). *J. Immunol.* 137, 245–254. *J. Immunol.* 186 (2011) 5024–5033.
- Elices, M.J., Osborn, L., Takada, Y., Crouse, C., Lohowskyj, S., Hemler, M.E., Lobb, R.R., 1990. VCAM-1 on activated endothelium interacts with the leukocyte integrin VLA-4 at a site distinct from the VLA-4/fibronectin binding site. *Cell* 60, 577–584.
- Frenette, P.S., Johnson, R.C., Hynes, R.O., Wagner, D.D., 1995. Platelets roll on stimulated endothelium in vivo: an interaction mediated by endothelial P-selectin. *Proc. Natl. Acad. Sci. U.S.A.* 92, 7450–7454.
- Hattori, R., Hamilton, K.K., Fugate, R.D., McEver, R.P., Sims, P.J., 1989. Stimulated secretion of endothelial von Willebrand factor is accompanied by rapid redistribution to the cell surface of the intracellular granule membrane protein GMP-140. *J. Biol. Chem.* 264, 7768–7771.
- Heja, D., Harmat, V., Fodor, K., Wilmanns, M., Dobo, J., Kekesi, K.A., Zavodszky, P., Gal, P., Pal, G., 2012a. Monospecific inhibitors show that both mannan-binding lectin-associated serine protease-1 (MASP-1) and -2 are essential for lectin

- pathway activation and reveal structural plasticity of MASP-2. *J. Biol. Chem.* 287, 20290–20300.
- Heja, D., Kocsis, A., Dobo, J., Szilagyi, K., Szasz, R., Zavodszky, P., Pal, G., Gal, P., 2012b. Revised mechanism of complement lectin-pathway activation revealing the role of serine protease MASP-1 as the exclusive activator of MASP-2. *Proc. Natl. Acad. Sci. U.S.A.* 109, 10498–10503.
- Hess, K., Ajjan, R., Phoenix, F., Dobo, J., Gal, P., Schroeder, V., 2012. Effects of MASP-1 of the complement system on activation of coagulation factors and plasma clot formation. *PLoS One* 7, e35690.
- Jani, P.K., Kajdacs, E., Megyeri, M., Dobo, J., Doleschall, Z., Futosi, K., Timar, C.I., Mocsa, A., Mako, V., Gal, P., Cervenak, L., 2014. MASP-1 induces a unique cytokine pattern in endothelial cells: a novel link between complement system and neutrophil granulocytes. *PLoS One* 9, e87104.
- Jenny, L., Dobo, J., Gal, P., Schroeder, V., 2015. MASP-1 of the complement system promotes clotting via prothrombin activation. *Mol. Immunol.* 65, 398–405.
- Kansas, G.S., 1996. Selectins and their ligands: current concepts and controversies. *Blood* 88, 3259–3287.
- Kiszel, P., Mako, V., Prohaszka, Z., Cervenak, L., 2007. Interleukin-6–174 promoter polymorphism does not influence IL-6 production after LPS and IL-1 beta stimulation in human umbilical cord vein endothelial cells. *Cytokine* 40, 17–22.
- Kjaer, T.R., Thiel, S., Andersen, G.R., 2013. Toward a structure-based comprehension of the lectin pathway of complement. *Mol. Immunol.* 56, 222–231.
- Kjaer, T.R., Le, T.M., Pedersen, J.S., Sander, B., Golas, M.M., Jensenius, J.C., Andersen, G.R., Thiel, S., 2015. Structural insights into the initiating complex of the lectin pathway of complement activation. *Structure* 23, 342–351.
- Korney, Z., Beke, S., Mihalfy, T., Jelitai, M., Kovacs, K.J., Szabo, Z., Szabo, B., 2013. Cell sorting in a Petri dish controlled by computer vision. *Sci. Rep.* 3, 1088.
- Krarup, A., Gulla, K.C., Gal, P., Hajela, K., Sim, R.B., 2008. The action of MBL-associated serine protease 1 (MASP1) on factor XIII and fibrinogen. *Biochim. Biophys. Acta* 1784, 1294–1300.
- Kubes, P., Kanwar, S., 1994. Histamine induces leukocyte rolling in post-capillary venules. A P-selectin-mediated event. *J. Immunol.* 152, 3570–3577.
- Lomakina, E.B., Waugh, R.E., 2009. Adhesion between human neutrophils and immobilized endothelial ligand vascular cell adhesion molecule 1: divalent ion effects. *Biophys. J.* 96, 276–284.
- Mako, V., Czucz, J., Weiszhar, Z., Herczenik, E., Matko, J., Prohaszka, Z., Cervenak, L., 2010. Proinflammatory activation pattern of human umbilical vein endothelial cells induced by IL-1beta, TNF-alpha, and LPS. *Cytometry A* 77, 962–970.
- Matsushita, M., Endo, Y., Fujita, T., 2013. Structural and functional overview of the lectin complement pathway: its molecular basis and physiological implication. *Arch. Immunol. Ther. Exp. (Warsz)* 61, 273–283.
- McLaughlin, F., Hayes, B.P., Horgan, C.M., Beesley, J.E., Campbell, C.J., Randi, A.M., 1998. Tumor necrosis factor (TNF)-alpha and interleukin (IL)-1beta down-regulate intercellular adhesion molecule (ICAM)-2 expression on the endothelium. *Cell Adhes. Commun.* 6, 381–400.
- Megyeri, M., Mako, V., Beinrohr, L., Doleschall, Z., Prohaszka, Z., Cervenak, L., Zavodszky, P., Gal, P., 2009. Complement protease MASP-1 activates human endothelial cells: PAR4 activation is a link between complement and endothelial function. *J. Immunol.* 183, 3409–3416.
- Megyeri, M., Harmat, V., Major, B., Vegh, A., Balczer, J., Heja, D., Szilagyi, K., Datz, D., Pal, G., Zavodszky, P., Gal, P., Dobo, J., 2013. Quantitative characterization of the activation steps of mannan-binding lectin (MBL)-associated serine proteases (MASPs) points to the central role of MASP-1 in the initiation of the complement lectin pathway. *J. Biol. Chem.* 288, 8922–8934.
- Megyeri, M., Jani, P.K., Kajdacs, E., Dobo, J., Schwaner, E., Major, B., Rigo Jr., J., Zavodszky, P., Thiel, S., Cervenak, L., Gal, P., 2014. Serum MASP-1 in complex with MBL activates endothelial cells. *Mol. Immunol.* 59, 39–45.
- Olofsson, A.M., Arfors, K.E., Ramezani, L., Wolitzky, B.A., Butcher, E.C., von Andrian, U.H., 1994. E-selectin mediates leukocyte rolling in interleukin-1-treated rabbit mesentery venules. *Blood* 84, 2749–2758.
- Oroszlan, M., Herczenik, E., Rugonfalvi-Kiss, S., Roos, A., Nauta, A.J., Daha, M.R., Gombos, I., Karadi, I., Romics, L., Prohaszka, Z., Fust, G., Cervenak, L., 2006. Proinflammatory changes in human umbilical cord vein endothelial cells can be induced neither by native nor by modified CRP. *Int. Immunol.* 18, 871–878.
- Petheo, G.L., Orient, A., Barath, M., Kovacs, I., Rethi, B., Lanyi, A., Rajki, A., Rajnavolgyi, E., Geiszt, M., 2010. Molecular and functional characterization of Hv1 proton channel in human granulocytes. *PLoS One* 5, e14081.
- Pietersma, A., Tilly, B.C., Gaestel, M., de Jong, N., Lee, J.C., Koster, J.F., Sluiter, W., 1997. p38 mitogen activated protein kinase regulates endothelial VCAM-1 expression at the post-transcriptional level. *Biochem. Biophys. Res. Commun.* 230, 44–48.
- Pober, J.S., Lapierre, L.A., Stolpen, A.H., Brock, T.A., Springer, T.A., Fiers, W., Bevilacqua, M.P., Mendrick, D.L., Gimbrone Jr., M.A., 1987. Activation of cultured human endothelial cells by recombinant lymphotoxin: comparison with tumor necrosis factor and interleukin 1 species. *J. Immunol.* 138, 3319–3324.
- Pudelko, M., Bull, J., Kunz, H., 2010. Chemical and chemoenzymatic synthesis of glycopeptide selectin ligands containing sialyl Lewis X structures. *ChemBiochem* 11, 904–930.
- Rondaij, M.G., Bierings, R., Kragt, A., van Mourik, J.A., Voorberg, J., 2006. Dynamics and plasticity of Weibel-Palade bodies in endothelial cells. *Arterioscler. Thromb. Vasc. Biol.* 26, 1002–1007.
- Salanki, R., Hos, C., Orgovan, N., Peter, B., Sandor, N., Bajtay, Z., Erdei, A., Horvath, R., Szabo, B., 2014. Single cell adhesion assay using computer controlled micropipette. *PLoS One* 9, e111450.
- Shimaoka, M., Lu, C., Palframan, R.T., von Andrian, U.H., McCormack, A., Takagi, J., Springer, T.A., 2001. Reversibly locking a protein fold in an active conformation with a disulfide bond: integrin alphaL I domains with high affinity and antagonist activity in vivo. *Proc. Natl. Acad. Sci. U.S.A.* 98, 6009–6014.
- Soh, U.J., Dores, M.R., Chen, B., Trejo, J., 2010. Signal transduction by protease-activated receptors. *Br. J. Pharmacol.* 160, 191–203.
- Thompson, P.W., Randi, A.M., Ridley, A.J., 2002. Intercellular adhesion molecule (ICAM)-1 but not ICAM-2, activates RhoA and stimulates c-fos and rhoA transcription in endothelial cells. *J. Immunol.* 169, 1007–1013.
- Tonnesen, M.G., 1989. Neutrophil-endothelial cell interactions: mechanisms of neutrophil adherence to vascular endothelium. *J. Invest. Dermatol.* 93, 53S–58S.
- Tucker, K.A., Lilly, M.B., Heck Jr., L., Rado, T.A., 1987. Characterization of a new human diploid myeloid leukemia cell line (PLB-985) with granulocytic and monocytic differentiating capacity. *Blood* 70, 372–378.
- Volanti, C., Gloire, G., Vanderplasschen, A., Jacobs, N., Habraken, Y., Piette, J., 2004. Downregulation of ICAM-1 and VCAM-1 expression in endothelial cells treated by photodynamic therapy. *Oncogene* 23, 8649–8658.
- Walz, G., Aruffo, A., Kolanus, W., Bevilacqua, M., Seed, B., 1990. Recognition by ELAM-1 of the sialyl-Lex determinant on myeloid and tumor cells. *Science* 250, 1132–1135.